

# *TRICKS OF THE TRADE*

L. Synn Stern  
*MPh, DV8-RTD*

# APPEARANCE

## *SAFETY*

- Shoes should come off easily or be appropriate for running in.
- Long earrings or big hoops for pierced ears may get pulled accidentally or intentionally. Wear small earrings, *clip-ons* or none at all
- Avoid necklaces, scarves. Across-the-body shoulder bags, or anything else that can accidentally or intentionally be tightened around your throat.
- Wigs should fit so that they can't slip and cover the face.
- Wear clothing that doesn't have to be removed to fuck. This saves time, hides money and tracks, and if there's a problem you won't lose your clothing if you have to run away or leave in a hurry.
- Clothing should be "straitjacket-proof." A half zipped or buttoned jacket can be pulled over shoulders to trap arms.
- Wear nothing that can get caught in or on car doors, like loose bracelets, billowing dresses. wide sleeves, or long coats with dangling, attached belts.
- Try to keep pubic hair clipped short - it won't get pulled either by accident or intentionally
- Make yourself noticeable/attractive to customers, and ensure that you can see them as well. If you need glasses or contact lenses, wear them- Constricted pupils do not admit too much light , so limit your heroin intake if you work at night.
- "Act as if." Always pretend to have security, wealth, health and calm.

## *HEALTH*

- Tight pants, nylon underwear and pantyhose can cause vaginal infections. Wear cotton or crotchless underwear. stockings, or nothing. Change panties daily (or turn inside out on alternate days, or use pantyliners). The same applies for socks.
- Keep toenails short to reduce toot pain. Shoes that are too large are safer and more comfortable than shoes that are too small.
- Sanitary napkins can provide a pathway for bacteria from anus to vagina, causing infection. Wear tampons while having your period and change them frequently.
- Keep lips, teeth, gums, vagina and thighs well lubricated to prevent irritation.
- Brush your teeth, douche and use enemas after work, not before or between customers,

# NEGOTIATION WITH CUSTOMERS

## ***BEFORE GOING TO WORK***

- Have a price list and stick to it
- Decide what you are and are not willing to do.
- Pick your own parking spot or hotel, don't leave it up to the customer
- Have a time limit for each service. If a customer can't come within a reasonable period he's costing you money by preventing you from seeing other customers. His failure to get off may make him agitated or violent
- Urinate often- friction from fucking with a full bladder may cause a bladder infection
- If you have access to "female" condoms check them out They can be used for both vaginal and anal sex, are controlled by the wearer, and they offer protection from both pregnancy and sexually transmitted diseases.
- Female condoms can be inserted before you go out to the stroll and will guarantee some protection.
- Have a supply of condoms, lube, and napkins with you and put a condom in your mouth
- Be willing to turn down particular customers, particular requests and offers of payment below your standards.
- Use some mentholated salve, like Vicks Vapo-Rub (for chapped lips & colds) inside your nostrils - you'll smell the customer less when you give head and you'll be less snifflly if you're dope sick.

## **REMEMBER**

- Run against traffic, not with it
- The smaller the stroll, the more you have to guard your reputation. In a small town with a small pool of customers, you can't afford to get caught stealing, faking, going with someone for less-than-price or without condoms.
- The further from the stroll you get with a customer, the harder it becomes to turn him down.
- If he gives you a bad feeling on the stroll (your territory) It will get worse once you're in the car (his territory).
- Stand where you can be seen and where you have good, glare-free vision of the stroll and approaching cars.
- Stash keys and other important personal items where they will be accessible and safe - keeping them in your pockets or purse makes them available to customers who want to rob you,

## ***APPROACHING THE CAR***

- Approach the driver's window keeping enough distance to avoid being grabbed, assaulted with spit or spray paint or pleasing "window shoppers"
- Study the customer, the car and the door lock system.
- Arrange price, service and location while you're outside the car.
- Circle the car completely before entering to ensure the number of passengers and to take the license plate number or pretend to.
- Wave goodbye to work partner (or pretend to) and shout expected return time.

## ***IN THE CAR***

- Ensure that you can re-open the car door.
- Get a good look at the customer and the car. Its still not too late to back out. For example, he's drunker than you thought you see a weapon under his seat or you get a creepy feeling.
- Get your money up front.
- Let him see you stash his payment and keep it separate from the rest of your money. If he tries to rob you maybe he won't find your entire bankroll.
- Put your purse where you can find it without looking. For example, open on the floor between your feet.
- Act in control. and take charge of the action.
- Keep eyes on him. and his hands, at all times.
- Don't let him get between you and the exit
- Keep the door open a crack if he allows.

# SERVICES

## *FELLATIO (Blow jobs)*

- The best positions in a car are in the passenger seat or crouching outside his door. The worst position is in the footwell crouching at his feet (this is usually only possible in vehicles that have lots of foot room) - you can be trapped there very easily. In a two-door vehicle, try to remain in the front seat
- Use hands to get him hard, keep him hard enough to get a condom on and prevent him from coming too soon if this means you'll have to give his money back.
- Get condom on (with hands or mouth) and check to make sure its on properly,
- Try not to rest any part of your face (chin, cheeks, forehead) on his balls or thighs. If he has sores, herpes, lice you may got them too.
- Take movement am from him.
- Breathe through your nose and don't close your eyes.
- If he wants to touch his dick and doesn't know he's wearing a condom distract him into touching you.
- To deep throat make a straight line from your chin to neck- This is difficult from the passenger seat unless you crouch on the seat
- If you don't like deep contact, tuck your chin to your chest. In this position there's a real limit to how far in he can get.
- Scrotum tightening signals imminent ejaculation.
- Most men thrust when they come. Arrange your chin so he's a comfortable depth in your mouth.
- Once he comes, keep firm pressure on condom with lips and hands to prevent semen leaking into your mouth.
- If you have gotten a condom on without him knowing it, sneak it off with far side hand (your left if you're in the passenger seat) and hide it in a baggie or napkin in your purse.

## *SEXUALLY TRANSMITTED DISEASES AND BLOWJOBS*

- Its possible to get sexually transmitted diseases (STDs) like gonorrhea, syphilis, chlamydia, herpes, warts and HIV from giving head to an infected person.
- Warts caused by human papilloma virus (HPV) have been linked to cervical cancer in women
- HPV is incurable, but warts can be surgically removed.
- HPV in the mouth can affect larynx and vocal chords. Surgery here is complicated.
- Men you're infected with an STD you are at higher risk for infection with HIV and hepatitis. Accessing STD testing and treatment regularly will reduce your risk for other, more deadly, diseases.

## *IF YOU MUST GIVE HEAD WITHOUT A CONDOM*

- Keep lips, mouth and throat as wet as possible. Don't swallow your spit.
- Try to suck and control moves, rather than holding mouth slack and letting him move into you
- Throat tissue is fragile, try to keep contact shallow.

- If you must deep throat, don't panic. Tension will make your throat tighten and tissues easier to tear.
- It is better to have him come on you than to have him thrusting in your mouth.
- The risk of infection is due more to fluids entering cuts or abrasions in your mouth, lips, throat,
- than swallowing pre-cum and semen. Stomach acids will probably destroy infectious agents so swallowing may be safer than holding semen in your mouth while you look for a place to spit it out.
- As soon as possible when its over, gargle with astringent mouthwash, hydrogen peroxide, or hard
- liquor -don't brush your teeth.

## ***VAGINAL SEX***

- Birth control is necessary for all female sexworkers. Weigh the costs and benefits of each method.
- It's possible to get pregnant before you begin menstruating, after what you thought was menopause, and anytime in between, even if your cycles are infrequent.
- Female condoms are good—if you can get used to them—since you control them. They can be inserted before you go out to work and can be used more than once (although they should be cleaned with warm, soapy water between customers). The drawbacks of female condoms are that they are expensive and do not prevent the customer's semen from coming into contact with the area around your vagina, leaving you at some risk for STDs. Find out if your local AIDS service or family planning agency distributes them for free or at a discount
- Male condoms are imperfect, especially if not used properly. Rape remains an occupational hazard, especially on strolls where sexwork is illegal. Your local harm reduction, AIDS service or family planning agency may distribute free condoms.
- The pill, implants and “prickpill” are effective but can cause candida and depression.
- The I.U.D. is effective, but the string helps carry infectious agents from the vagina into uterus and tubes causing Pelvic Inflammatory Disease (PID) which is serious and sometimes life-threatening.
- Diaphragms must be used with spermicide and held in place for 8 hours after his last ejaculation- The benefits are that man can't feel it with the penis and its a cheap, reusable tampon. The problems are that it sometimes causes bladder infections, can cause infections if left in place too long and it must be refitted with changes in body weight.
- The cervical cap works like a diaphragm but is made from hard plastic, which men can feel with their penises.
- Spermicide foams, creams, jellies, sponges, suppositories and films are only effective when used with a condom. Nonoxynol 9, the most common spermicidal ingredient, may help kill HIV and some other STDs. However, these substances can be messy, drippy and can be irritating to vaginal, anal and penile tissue, especially when used daily.

## ***RISKS OF VAGINAL SEX***

- Pregnancy, sexually transmitted disease, HIV and hepatitis if you don't use latex barriers (male condoms, female condoms, dental dams).
- Bladder infections from friction of the penis over full bladder which, if untreated, can cause serious kidney infections

- Candida, trichomonads, hemophilus gardnerella from poor hygiene and from wearing tight clothing or synthetic fabrics over damp wet genitals.
- Irritation of labia from pulled hairs and local trauma
- Gas or difficulty shitting following trauma (fucking too hard).

### *RISK PREVENTION*

- Use condoms and appropriate birth control.
- Keep labia and inner thighs well lubricated with water soluble lubricant
- Re-activate lubricant with spit as needed
- Urinate frequently - ideally before and after every fuck, Some drugs (cocaine. speed) make it feel as if you don't need to piss. Do it anyway, and use hands on abdomen to press last drops of urine from bladder.
- Wipe front to back after pissing and shitting
- Wear crotchless underwear, clean cotton ones or none at all. Wear stockings instead of pantyhose.
- Drink lots of unsweetened fruit juices (citrus, cranberry) or take vitamin C to prevent bladder and yeast infections.
- Don't douche. If you must do it after work, not before, with a mild salt solution or white vinegar in warm water.
- Yogurt or a whole, unpeeled garlic clove in the vagina will kill candida (yeast)- Eating yogurt helps prevent candida.
- Too much sugar. coffee, tea, cola, cocaine. or speed will increase your risk of bladder or vaginal infections.
- Shower or bathe daily if possible. Avoid perfumed soap and bubble bath which can cause irritation and bladder infections.
- If you can't bathe, wash your genitals and allow them to dry completely before dressing
- Infections cause small cuts in mucous tissue which increase your chances of getting STDs, HIV and hepatitis.
- Prevent as much as you can- If you do get an infection get it treated immediately.

### *RISK REDUCTION WITH THE CUSTOMER*

- Try to negotiate out of performing vaginal or anal sex. Offer to do it with hands, mouth. between thighs, tits and ass cheeks

OR

- Charge a very high price for this service.
- Use a lubricated condom and lubricate entire genital area.
- Put condom on with your mouth if he won't permit one, and swing yourself on top of him quickly (while hiding his cock in your hands) so he doesn't see it
- In a two-door car, try to remain in the front seat.
- Position yourself on top and facing him. This way you can move easily and keep an eye on his hands.
- Open labia yourself after first spitting on fingers to reactivate lubricant and push hairs to the sides.
- Hold your pelvis so that the head of his penis is pointing to the middle of your basic This is the least friction-producing position for most women.
- The closer together your thighs are, the shallower he'll be in your vagina.

- When you're fucking someone who excites you, your vagina lubricates naturally, expands, and your cervix moves higher in your body- If this doesn't happen you may have some pain from the penis banging into your cervix
- If you have A LOT of pain when he bangs your cervix, you probably have an infection. See a medical provider as soon as possible.
- Never let him (or anyone else) put fingers first in your ass and then in your vagina - this will place you at high risk for infections.
- If you have bad cramps after fucking, check the position of your cervix with a clean finger. If you can't feel the os (hole) then the cervix got banged out of position. Try to reach behind it and pull it back into position. It helps to squat and push down as if you're shitting.
- Keep a hand on the base of his penis to:
  - 1) keep him hard
  - 2) make sure the condom doesn't roll up
  - 3) keep your labia spread and reduce trauma
  - 4) 4) limit depth of penis in your body
- Once he comes get him out fast or hold condom tight to the base of his penis so semen doesn't leak into you. If you snuck condom on, sneak it off. Don't spill semen on or near your vagina.
- Its hard to sneak condoms on without giving head. If it isn't possible to sneak one on, try appealing to
  - 1) his safety
  - 2) his libido (condoms make it last longer, condoms excite you and you'll fuck longer)
  - 3) his status as a customer - he's paying you as an expert in giving pleasure - he should trust that you'll be sure to please him while keeping both of you safe
  - 4) his sense of honor about your safety.
- If you find you're fucking a lot of customers without condoms, consider using a diaphragm. Some protection is better than none and this method offers partial protection against pregnancy and disease.

## ***ANAL SEX***

- Anal sex can be uncomfortable, make shitting more difficult and is a very efficient way of getting HIV, hepatitis B. and other diseases.
- Try to negotiate out of it with another service or charge so much that a cheap customer won't be willing to pay.
- If you think you can pull it off, try faking by using your hands, ass cheeks or vagina.
- Use a well lubricated thick condom and lots of extra lubricant on your anus
- Female condoms can be used for anal intercourse
- To prevent infections, don't let him go from your ass to your mouth or vagina. Forbid fingering or **insist** he use separate hands for the anus and vagina. If you must perform another service following anal sex, change condoms.
- Position yourself on top and facing him as for vaginal sex, but this time put your feet on the seat on either side of him (like a jockey on a horse). This straightens your colon so you will have less discomfort and you will be able to move more easily.
- When he enters, exhale slowly while pushing down like you're taking a shit (you won't). His penis will go in more easily,



- If you're nervous, your sphincter will tighten and can tear. Try to stay relaxed and control the movements.
- Keep a hand at the base of his dick to limit the depth and check that the condom doesn't roll up.
- If you have shit in your ass, it'll get pushed higher into your body when assfucking. You may have difficulty shitting the next day. You can prevent/treat constipation by drinking lots of water and/or fruit juice, eating fruits and vegetables or something that's oily/fatty.
- If constipated drink lots of water, eat some oily foods and/or fruit and vegetables, using coke or speed may help or use a reliable laxative pill. Or put a clean finger inside your vagina and move the shit down toward your anus and out your body through the membrane wall.
- If you must use an enema. do it an a day when you don't have to work.
- You might drip lubricant and a little loose shit for awhile- It can cause an infection if it gets in your vagina.
- Wipe front to back and if you want to catch the drip in a pantyliner, place the liner to the back so the bacteria can't "walk" along the pad and into your vagina and/or urethra.

### *ANALINGUS (RIMMING)*

- Fake it with spit on a knuckle while lowering your head- Wash your hands afterwards.
- If your can't fake it use a dental dam, a cut condom or latex glove, or a doubled up piece of plastic wrap.
- Using lube between your customer's asshole and the barrier will increase his sensation
- It's possible to get diseases like hepatitis A or B, worms and amoebas from licking the ass of a person who's infected.

### *CUNNILINGUS (EATING PUSSY)*

- Its risky to lay under a stranger- Stay on top.
- If he bites you it will hurt and you may get an infection. Even a minor irritation or trauma increases the chance that you'll get an infection during your next sexual encounter.
- If you have HFV or another STD you may pass it to somebody else this way, even if you're not menstruating.
- Use a dental dam, cut condom or latex glove, or a double layer of sandwich wrap.

### *MILKMAIDS*

- If you have HIV, it is possible to paw it to somebody else by letting them drink your breastmilk.
- It probably won't negatively affect your milk supply.

### *WATERSPORTS*

- Try to be the pisser, not the one who gets pissed on- Piss stinks, and your ability to attract another customer may be limited.
- Its probably safe to get urine in your mouth.

### *SCAT*

- It's never safe to get shit in your mouth (even your own shit), vagina or near your urethra•

- If you must eat his shit, seek medical advice afterwards. It might be a good idea to take antibiotics.
- If he wants to eat your shit and you can do it: go ahead. But don't let him lick your vagina afterwards.

### *S&M*

- Its never safe to let a stranger tie you up or hit you.
- Be the "top"
- Use rubber gloves (and diaphragm jelly on your hands for extra protection) if you plan to enter his anus, or there's a chance you'll draw blood.

## **SEXWORK AND PREGNANCY**

- Pregnancy may be “immunosuppressive” (your immune system may be weakened). Take extra care.
- Try selling only hand and mouth work.
- Go easy with vaginal sex. Use a condom, lubricant and avoid deep contact. Stay in control of the movements.
- Be alert for signs of infection, and treat immediately.
- You may develop hemorrhoids. These can bleed, itch, and provide another route for disease to enter your body.

## **SEXWORK AND DRUG USE**

- Try not to let customers know that you use or they may be tempted to play money and power games.
- Insist on getting paid in cash, not drugs - you'll look more professional and in control.
- Prevent track marks by rotating injection sites, using new, sharp, small gauge needles, and salve or hemorrhoid cream on injection sites and old tracks.
- Don't get so pinned that you can't see at night (use cocaine or speed, less dope. or substitute demerol)
- It's more difficult to work safe when you're very high on cocaine or other drugs.
- Try not to use alcohol at all when working - your reaction time will be slower.
- Keep mouth moist by chewing gum or sucking hard candies.
- Drink a lot of water to prevent constipation, dry mouth and dry, itchy skin.
- Don't *let* customers know that you're dope sick. Prevent sniffing with antihistamines and mentholated salve in your nostrils.
- Urinate frequently, even if you think you don't have to.

## **SELF DEFENSE**

- Like diseases, it is better and easier to prevent violence than to treat it. The best defense is a good offense. Be aware.
- Do not carry a weapon that can be taken from and used against you.
- Even small men are often stronger than large women. Do not try to fight a man unless you're certain that you can escape or that you will win..
- Try using your voice and speed instead of your strength- Make a lot of noise. Scream. Turn on the car lights. Hit the horn- Try to get the door open, attack and run.
- Attack areas that are easily injured. such as throat, eyes, adam's apple, nose, underside of nose, testicles, shins, insteps.
- Kick, poke with fingers, nails, keys, spray hairspray in his face
- Don't bother trying to pull his hands off your throat since he will probably be stronger. Break a
- strangle hold by joining hands. and swinging your arms up against him while simultaneously moving your body down and away.
- Run against traffic and towards lights and people.

## WHAT YOUR AGENCY CAN DO

- Provide condoms (male & female). lubricant dental dams and tampons,
- Make mouthwash, hemorrhoid cream, body lotion and sunscreen available.
- Provide new injection equipment (or bleach kits) and alcohol wipes.
- Offer the use of a mirror, toilet shower, telephone, washing machine and dryer.
- Post "ugly-mug" sheets and keep them up-to-date.
- Report rapes and assaults.
- Know the local prostitution laws
- Report abusive police.
- Offer drug and financial advisory services, and referral to nonjudgmental providers of medical. legal and other professional services
- Display clothing and accessories appropriate to safe street sex work. Keep a supply of donated clothing and umbrellas on hand.
- Provide vitamin C rich foods such as tomato soup and citrus and cranberry juices to prevent bladder infections, bleeding gums and easy bruising.
- Encourage sexworkers to talk to work with, and support each other.
- Discourage sexworkers from undercutting each others' prices.
- Teach self-defense, meditation, relaxation and creative visualization techniques.
- Help sexworkers develop good street skills, especially intuition/instinct
- Understand that superstition has a role in any unsafe or unpredictable profession or situation. Help sexworkers develop superstitions that are actually protective. Do this by helping them ADD to their superstition pool, not by trying to convince them that their current beliefs are invalid.
- Support sexworkers who want to leave the sex industry as well as those who continue to work in it
- Cosmetics, hygiene supplies, cotton underwear, and stockings make nice gifts.